

Regularization of Ill-Posed Problems with Unbounded Operators. ^{*}†

A.G. Ramm

Mathematics Department, Kansas State University,
Manhattan, KS 66506-2602, USA
ramm@math.ksu.edu

Abstract

Variational regularization and the quasisolutions method are justified for unbounded closed operators.

1 Introduction. Variational regularization.

There is a large literature on methods for solving ill-posed problems: variational regularization, method of quasisolutions, iterative and projection regularization [2]-[8]. The case of ill-posed problems with a closed linear operator was discussed in [5], and the case of nonlinear, possibly unbounded, operators possibly unbounded, does not seem to be discussed. In the theory of ill-posed problems the following well-known result ([1, Lemma I.5.8]) is often used: if A is an injective and continuous mapping from a compact set M of a Banach space into a set $N := AM$, then the inverse mapping $A^{-1} : N \rightarrow M$ is continuous. In [7, p.112] the usual assumption about continuity of A in the above result is replaced by the assumption about closedness of A .

In this short note ill-posed problems are studied in the case of the mapping A not necessarily continuous, but closed, possibly nonlinear. Our argument is very simple and the result is fairly general.

Let A be an injective, possibly nonlinear, closed operator on a Banach space X , and the equation

$$Ay = f \tag{1.1}$$

has a solution y . Our arguments hold in metric spaces as well without changes.

Assume that A^{-1} is not continuous. This implies that problem (1.1) is ill-posed. Let

$$\|f_\delta - f\| \leq \delta. \tag{1.2}$$

^{*}key words: regularization, ill posed problems, unbounded operators

[†]Math subject classification: 47A52 65J 20

Given f_δ and A , one wants to construct $u_\delta = R_\delta(f_\delta)$, such that $\|u_\delta - y\| \rightarrow 0$ as $\delta \rightarrow 0$, where y solves (1.1). If u_δ is constructed, then the operator R_δ yields a stable approximation of the solution y to (1.1).

Let us first describe the method of variational regularization in our case.

Define the functional

$$F(u) := \|A(u) - f_\delta\| + \delta\phi(u), \quad (1.3)$$

and assume that $\phi(u) \geq 0$ is a functional, such that for any constant $c > 0$ the set

$$\{u : \phi(u) \leq c\} \text{ is precompact in } X. \quad (1.4)$$

The functional F depends on δ and f_δ , but for simplicity of writing we do not show this dependence explicitly. Assume that $D(A) \subset D(\phi)$, the domain of definition of ϕ , contains $D(A)$. This assumption implies that $y \in D(\phi)$, so that $\phi(y) < \infty$. Define $D(F) = D(A)$. If A were bounded, defined on all of X , then one would assume $y \in D(\phi)$ and $D(F) = D(\phi)$. If A were unbounded and $D(\phi) \subset D(A)$, then one would assume that $y \in D(\phi)$ and $D(F) = D(\phi)$.

Denote

$$0 \leq m := \inf_{u \in D(A)} F(u). \quad (1.5)$$

The number $m = m(\delta) \geq 0$. Let u_j be a minimizing sequence $u_j \in D(F)$ for the functional F , such that:

$$F(u_j) \leq m + \frac{1}{j} = m + \delta, \quad \frac{1}{j} \leq \delta. \quad (1.6)$$

Denote by $u_\delta := u_{j(\delta)}$ a member $u_{j(\delta)}$ of this minimizing sequence, where $j(\delta)$ is chosen so that $\frac{1}{j(\delta)} \leq \delta$. There are many such $j(\delta)$ and we fix one of them, for example, the minimal one. Since

$$F(y) \leq \delta + \delta\phi(y) := c_1\delta, \quad c_1 := 1 + \phi(y), \quad (1.7)$$

one has:

$$m \leq c_1\delta, \quad (1.8)$$

and

$$F(u_\delta) \leq m + \delta \leq c\delta, \quad c := c_1 + 1. \quad (1.9)$$

Thus $\delta\phi(u_\delta) \leq c\delta$, and

$$\phi(u_\delta) \leq c. \quad (1.10)$$

Let us now take $\delta \rightarrow 0$. By (1.4) and (1.10) one can select a convergent in X subsequence of the set u_δ , which we denote also u_δ , such that

$$\|u_\delta - u\| \rightarrow 0 \text{ as } \delta \rightarrow 0, \quad (1.11)$$

where u is the limit of u_δ .

From (1.2), (1.3), (1.9) and (1.10), one gets

$$0 = \lim_{\delta \rightarrow 0} F(u_\delta) = \lim_{\delta \rightarrow 0} \|A(u_\delta) - f_\delta\| = \lim_{\delta \rightarrow 0} \|A(u_\delta) - f\|. \quad (1.12)$$

Since A is closed, (1.11) and (1.12) imply

$$\lim_{\delta \rightarrow 0} A(u_\delta) = A(u), \quad 0 = \|A(u) - f\|. \quad (1.13)$$

Since A is injective, (1.13) and (1.1) imply $u = y$, so

$$\lim_{\delta \rightarrow 0} \|u_\delta - y\| = 0. \quad (1.14)$$

Since the limit y of any subsequence u_δ is unique, the whole sequence u_δ converges to y .

We have proved the following result:

Theorem 1.1. *Assume that (1.4) holds, $\phi \geq 0$, $A : D(A) \rightarrow X$ is a closed, injective, possibly nonlinear unbounded operator, $A(y) = f$, and A^{-1} is not continuous. Let u_δ be constructed as above so that (1.9) holds. Then (1.14) holds.*

In section 2 the method of quasisolutions is discussed in the case of possibly unbounded and nonlinear operators.

2 Quasisolutions for unbounded operators.

In this section the assumptions about equation (1.1) and the operator A are the same as in section 1, in particular, A^{-1} is not continuous, so that solving equation (1.1) is an ill-posed problem.

Choose a compactum $K \subset X$ such that the solution of (1.1) $y \in K$. Consider the problem

$$\|A(u) - f_\delta\| = \inf := \mu, \quad u \in K \subset D(A). \quad (2.1)$$

The infimum $\mu = \mu(\delta) \geq 0$ depends on f_δ also, but we do not show this dependence explicitly. Let u_j be a minimizing sequence:

$$\|A(u_j) - f_\delta\| \leq \mu + \frac{1}{j}. \quad (2.2)$$

Choose $j = j(\delta)$ such that $\frac{1}{j} \leq \delta$ and denote $u_j := u_\delta$.

Then

$$\|A(u_\delta) - f_\delta\| \leq \mu + \delta. \quad (2.3)$$

Since $\|A(y) - f_\delta\| \leq \delta$, it follows that $\mu \leq \delta$, so

$$\|A(u_\delta) - f_\delta\| \leq 2\delta. \quad (2.4)$$

Since $\{u_\delta\} \subset K$ one can select a convergent (to some u) subsequence, denoted also $\{u_\delta\}$:

$$\|u_\delta - u\| \rightarrow 0 \text{ as } \delta \rightarrow 0. \quad (2.5)$$

From (2.4) it follows that $A(u_\delta)$ converges to f :

$$\|A(u_\delta) - f\| \leq \|A(u_\delta) - f_\delta\| + \|f_\delta - f\| \leq 3\delta \rightarrow 0 \text{ as } \delta \rightarrow 0. \quad (2.6)$$

Since A is closed, it follows from (2.5) and (2.6) that

$$A(u) = f. \quad (2.7)$$

Injectivity of A , equation (2.7), and the equation $A(y) = f$ imply $u = y$. We have proved:

Theorem 2.1. *Assume that $A : D(A) \rightarrow X$ is a closed, injective, possibly nonlinear and unbounded, operator, $A(y) = f$, (1.2) holds, K is a compact set in X , and $y \in K$. If $u_\delta \in K$ satisfies (2.4), then (1.14) holds.*

Acknowledgement. The author thanks Professor A.Yagola for useful discussions.

References

- [1] Dunford N., Schwartz J., Linear operators, General theory, Interscience Publishers, New York, 1958.
- [2] Engl H., Hanke M., Neubauer A., Regularization of inverse problems, Kluwer, Dordrecht, 1996.
- [3] Groetsch C., Inverse problems in mathematical sciences, Vieweg, Braunschweig, 1993.
- [4] Ivanov, V., Vasin, V., Tanana, V., Theory of linear ill-posed problems and applications, Nauka, 1978.
- [5] Liskovetz O., regularization of equations with a closed linear operator, Diff. Equations, 7, N9, (1970), 972-976.
- [6] Morozov V., Methods of solving incorrectly posed problems, Springer Verlag, New York, 1984.
- [7] Ramm, A.G., Random fields estimation theory, Longman Scientific and Wiley, New York, 1990.
- [8] Tikhonov A., Leonov, A., Yagola, A., Nonlinear ill-posed problems, vol. 1,2, Chapman and Hall, London, 1998.